



University of
Salford
MANCHESTER

Genomic characterisation of the indigenous Irish Kerry cattle breed

Browett, S, McHugo, G, Richardson, IW, Magee, DA, Park, SDE, Fahey, AG, Kearney, JF, Correia, CN, Randhawa, IAS and MacHugh, DE

<http://dx.doi.org/10.3389/fgene.2018.00051>

| | |
|-----------------------|---|
| Title | Genomic characterisation of the indigenous Irish Kerry cattle breed |
| Authors | Browett, S, McHugo, G, Richardson, IW, Magee, DA, Park, SDE, Fahey, AG, Kearney, JF, Correia, CN, Randhawa, IAS and MacHugh, DE |
| Type | Article |
| URL | This version is available at: http://usir.salford.ac.uk/46300/ |
| Published Date | 2018 |

USIR is a digital collection of the research output of the University of Salford. Where copyright permits, full text material held in the repository is made freely available online and can be read, downloaded and copied for non-commercial private study or research purposes. Please check the manuscript for any further copyright restrictions.

For more information, including our policy and submission procedure, please contact the Repository Team at: usir@salford.ac.uk.



Genomic Characterisation of the Indigenous Irish Kerry Cattle Breed

Sam Browett¹, Gillian McHugo², Ian W. Richardson³, David A. Magee², Stephen D. E. Park³, Alan G. Fahey², John F. Kearney⁴, Carolina N. Correia², Imtiaz A. S. Randhawa⁵ and David E. MacHugh^{2,6*}

¹ Ecosystems and Environment Research Centre, School of Environment and Life Sciences, University of Salford, Salford, United Kingdom, ² Animal Genomics Laboratory, UCD School of Agriculture and Food Science, University College Dublin, Dublin, Ireland, ³ IdentiGEN Ltd., Dublin, Ireland, ⁴ Irish Cattle Breeding Federation, Bandon, Ireland, ⁵ Sydney School of Veterinary Science, The University of Sydney, Camden, NSW, Australia, ⁶ UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Dublin, Ireland

OPEN ACCESS

Edited by:

Ino Curik,

Faculty of Agriculture, University of Zagreb, Croatia

Reviewed by:

Gábor Mészáros,

University of Natural Resources and Life Sciences, Vienna, Austria

Paolo Ajmone Marsan,

Università Cattolica del Sacro Cuore, Italy

*Correspondence:

David E. MacHugh
david.machugh@ucd.ie

Specialty section:

This article was submitted to
Livestock Genomics,
a section of the journal
Frontiers in Genetics

Received: 03 September 2017

Accepted: 02 February 2018

Published: 19 February 2018

Citation:

Browett S, McHugo G, Richardson IW, Magee DA, Park SDE, Fahey AG, Kearney JF, Correia CN, Randhawa IAS and MacHugh DE (2018) Genomic Characterisation of the Indigenous Irish Kerry Cattle Breed. *Front. Genet.* 9:51.
doi: 10.3389/fgene.2018.00051

Kerry cattle are an endangered landrace heritage breed of cultural importance to Ireland. In the present study we have used genome-wide SNP array data to evaluate genomic diversity within the Kerry population and between Kerry cattle and other European breeds. Patterns of genetic differentiation and gene flow among breeds using phylogenetic trees with ancestry graphs highlighted historical gene flow from the British Shorthorn breed into the ancestral population of modern Kerry cattle. Principal component analysis (PCA) and genetic clustering emphasised the genetic distinctiveness of Kerry cattle relative to comparator British and European cattle breeds. Modelling of genetic effective population size (N_e) revealed a demographic trend of diminishing N_e over time and that recent estimated N_e values for the Kerry breed may be less than the threshold for sustainable genetic conservation. In addition, analysis of genome-wide autozygosity (F_{ROH}) showed that genomic inbreeding has increased significantly during the 20 years between 1992 and 2012. Finally, signatures of selection revealed genomic regions subject to natural and artificial selection as Kerry cattle adapted to the climate, physical geography and agro-ecology of southwest Ireland.

Keywords: cattle, conservation genomics, endangered breed, inbreeding, genetic diversity, population genomics, selection signature, single nucleotide polymorphism

INTRODUCTION

Approximately 10,000 years ago, humans first domesticated wild aurochs (*Bos primigenius*)—the progenitor of modern cattle—in the Fertile Crescent region of Southwest Asia (Larson and Fuller, 2014; Larson et al., 2014; MacHugh et al., 2017). Extant domestic cattle, which encompass humpless taurine (*B. taurus*), humped zebu (*B. indicus*) and myriad *B. taurus/indicus* hybrid populations, have, through genetic drift and natural and artificial selection, diversified into more than 1,100 recognised breeds. However, beginning in the middle of the twentieth century, socioeconomic preferences for large highly productive dairy, beef and dual-purpose breeds have led to extinction and increased vulnerability of more than 200 locally-adapted landrace or native cattle breeds (Gandini et al., 2004; Food and Agriculture Organization, 2007, 2015).

With the advent of accelerating climate change, particularly in the Arctic and circumarctic regions (Vihma, 2014; Gao et al., 2015), agro-ecological environments in north-western Europe

will inevitably undergo significant change during the coming century (Smith and Gregory, 2013; Wheeler and von Braun, 2013). It is, therefore, increasingly recognised that long-term sustainability of animal production systems and food security will necessitate conservation and management of livestock genetic resources in this region (Hoffmann, 2010; Boettcher et al., 2015; Kantanen et al., 2015). Locally-adapted native livestock breeds with distinct microevolutionary histories and minimal external gene flow will have accumulated novel genomic variation and haplotype combinations for quantitative health, fertility and production traits (Hill, 2014; Feliuss et al., 2015; Kristensen et al., 2015). These populations may therefore be key to future breeding programmes directed towards adaptation of European livestock to new agro-ecological and production environments (Biscarini et al., 2015; Boettcher et al., 2015; Phocas et al., 2016a,b).

The availability of powerful and cheap tools for genotyping large numbers of single nucleotide polymorphisms (SNPs) has provided conservation biologists and animal geneticists with the opportunity to characterise genomic variation and estimate population genetic parameters at very high resolution in threatened or endangered livestock breeds (Pertoldi et al., 2014; Ben Jemaa et al., 2015; Beynon et al., 2015; Mészáros et al., 2015; Burren et al., 2016; Decker et al., 2016; Iso-Touru et al., 2016; Manunza et al., 2016; Mastrangelo et al., 2016; Visser et al., 2016; Williams et al., 2016; François et al., 2017). These studies are already providing important baseline data for genetic conservation and will underpin programmes for managed breeding and biobanking of these populations (Groeneveld et al., 2016).

As a native breed with a claimed ancient heritage, Kerry cattle are considered culturally important to Ireland (Curran, 1990). It is a landrace cattle population that remains productive in harsh upland regions with poor quality feed, which are typical of southwest Ireland where the Kerry breed evolved (Food and Agriculture Organization, 2017). These cattle were often referred to anecdotally in Ireland as the “poor man’s cow” due to their ability to produce relatively large quantities of milk on very sparse fodder; the Kerry breed is also considered to be a remnant of what was once a substantially larger and more widespread historical population. Levels of inbreeding have been estimated using pedigree data and the accumulated figure since the foundation of the herd book in 1887 reached 15% in 1985 (O’hUigín and Cunningham, 1990).

In recent decades the Kerry cattle breed has experienced significant population fluctuations due to changing socioeconomic and agricultural circumstances. During the 1980s, the number of breeding females decreased to less than 200, prompting the Irish agricultural authorities to introduce a Kerry cattle conservation scheme (McParland, 2013), which has continued to the present day in the form of the Department of Agriculture, Food and the Marine (DAFM) Kerry Cattle Premium Scheme (Department of Agriculture Food and the Marine, 2017).

The formal conservation policy and supports initiated during the early 1990s led to a significant increase in the Kerry cattle population, such that by 2007 the number of breeding females had increased to more than a thousand animals (Food and

Agriculture Organization, 2007). In recent years, however, due to deteriorating economic circumstances in Ireland post-2008, the Kerry cattle population has substantially declined once again and is classified as endangered and under significant threat of extinction or absorption through crossbreeding with other breeds (McParland, 2013; Department of Agriculture Food and the Marine, 2014).

The Kerry cattle breed was one of the first European heritage cattle breeds to be surveyed using molecular population genetics techniques. We have previously used autosomal microsatellite genetic markers and mitochondrial DNA (mtDNA) control region sequence variation for comparative evolutionary studies of genetic diversity in Kerry cattle and other British, European, African and Asian breeds (MacHugh et al., 1997, 1998, 1999; Troy et al., 2001). In addition, Bray et al. have used microsatellites to examine admixture and ancestry in Kerry cattle and the Dexter and Devon breeds (Bray et al., 2009). Results from these studies demonstrated that Kerry cattle exhibit markedly low mtDNA sequence diversity, but autosomal microsatellite diversity comparable to other cattle breeds native to Britain and Ireland. More recently, analyses of medium- and high-density SNP genotypes generated using genome sequence data from an extinct British *B. primigenius* subfossil have shown that Kerry cattle retain a significant genomic signature of admixture from wild aurochs (Park et al., 2015; Upadhyay et al., 2017). This observation highlights the genetic distinctiveness of the Kerry population and has major implications for conservation and management of the breed.


For the present study, and within a genetic conservation framework, we performed high-resolution comparative population genomics analyses of Kerry cattle and a range of British and European cattle breeds. These analyses encompassed phylogenetic network reconstruction, evaluation of genetic structure and inbreeding, modelling of historical effective population sizes and functional analyses of artificial and natural selection across the Kerry genome.

MATERIALS AND METHODS

Kerry Cattle Population DNA Sampling in 1991/92 and 2011/12

Two different population samples from the Irish Kerry cattle breed were used for this study (Figure 1). The first population sample consisted of peripheral blood and semen straw genomic DNA collected and purified from 36 male and female Kerry cattle in 1991/92, which are a subset of the Kerry cattle population sample ($n = 40$) we have previously described and used for microsatellite-based population genetics analyses (MacHugh et al., 1997, 1998). Pedigree records and owners were consulted to ensure that a representative sample of animals was obtained. This Kerry population sample group is coded as KY92.

The second Kerry cattle population sample was collected in 2011/12 from 19 different herds located across southern and western Ireland. Performagene (PG-100) nasal swab DNA collection kits were used for biological sample collection (DNA



Genotek Inc., Ottawa, Canada). Nasal swab DNA samples were collected from a total of 75 male and female Kerry cattle and owners were consulted to ensure that a representative sample of animals was obtained. This Kerry population sample group is coded as KY12. Genomic DNA was purified from 0.5 ml of each PG-100 nasal swab sample using the laboratory protocol recommended by the manufacturer (DNA Genotek Inc.).

SNP Genotyping and Assembly of Comparative SNP Data Sets

Illumina® Bovine SNP50 BeadChip (Matukumalli et al., 2009) genotyping on all 111 Kerry genomic DNA samples (KY92 and KY12 sample panels plus nine blinded sample duplicates for quality control purposes) was performed by Weatherbys Scientific (Co. Kildare, Ireland).

For comparative population genomics analyses, equivalent SNP data for a range of other breeds were obtained from previously published work (Decker et al., 2009; Flori et al., 2009; Gibbs et al., 2009; Matukumalli et al., 2009; Gautier et al., 2010; Park et al., 2015). The breed SNP data were split into two discrete composite data sets: a European breed SNP data set (EU) and a SNP data set for a subset of European breeds originating from Britain and Ireland (BI). A population sample of West African N'Dama *B. taurus* cattle from Guinea (NDAM) was also used as an outgroup for the phylogenetic analyses. **Table 1** provides detailed biogeographical information on the cattle breed samples used for the present study.

Sample Removal and Quality Control and Filtering of SNPs

Genomic non-exclusion—in other words, genome-wide SNP profiles completely compatible with parent-offspring relationship (with allowance for very low-level genotyping error)—were used to identify animals from the KY92 and KY12 population samples that were parent-offspring pairs. One of the two animals in each pair was then randomly removed to generate the working SNP data set. Following this procedure, quality control and filtering based on recorded SNP genotypes was performed as detailed below for the EU and BI data sets.

Prior to quality control and filtering there were 54,057 SNPs in the EU data set (608 animals, including KY92 and KY12) and in the BI data set (354 animals, including KY92 and KY12). SNP quality filtering was performed using PLINK version 1.07 (Purcell et al., 2007), such that individual SNPs with more than 10% missing data and a minor allele frequency (MAF) of ≤ 0.01 (1%) were removed from both data sets; however, for analyses of genomic inbreeding and runs of homozygosity (ROH) the MAF filtering threshold was not imposed. Only autosomal SNPs were retained and individual animal samples with a SNP call rate less than 90% were also removed from each of the two data sets.

SNP quality control and filtering were performed across breeds/populations (by data set) for construction of phylogenies and ancestor graphs, multivariate analysis, investigation of population structure and detection of signatures of selection. For intrapopulation analyses of effective population size (N_e)

TABLE 1 | Cattle breed/population samples used for the present study.

| Breed/population | Code | Data set | Breed purpose | Country of origin | Source |
|-----------------------|------|----------|--------------------|----------------------|---------|
| Angus | ANGU | BI/EU | Beef | Scotland | 1, 2 |
| Belted Galloway | BGAL | BI | Beef | Scotland | 2 |
| British Shorthorn | BSHN | BI/EU | Dual purpose | England | 2 |
| Brown Swiss | BRSW | EU | Dairy | Switzerland | 1, 2, 3 |
| Charolais | CHAR | EU | Beef | France | 1, 2, 3 |
| Devon | DEVN | BI | Beef | England | 2 |
| Dexter | DXTR | BI | Dual purpose | Ireland | 2 |
| English Longhorn | ELHN | BI | Beef | England | 2 |
| Finnish Ayrshire | FAYR | BI/EU | Dairy | Scotland/Finland | 2 |
| Galloway | GALL | BI | Beef | Scotland | 2 |
| Gelbvieh | GELB | EU | Dual purpose | Germany | 2, 4 |
| Guernsey | GNSY | BI/EU | Dairy | Channel Islands | 1, 2 |
| Hereford | HRFD | BI/EU | Beef | England | 1, 2 |
| Holstein | HOLS | EU | Dairy | The Netherlands | 1, 2, 5 |
| Jersey | JRSY | BI/EU | Dairy | Channel Islands | 1, 2, 3 |
| Kerry sampled 1991/92 | KY92 | BI/EU | Dairy | Ireland | Current |
| Kerry sampled 2011/12 | KY12 | BI/EU | Dairy | Ireland | Current |
| Limousin | LIMS | EU | Beef/draft | France | 1, 2 |
| Lincoln Red | LNCR | BI | Beef | England | 2 |
| Montbeliarde | MONT | EU | Dairy | France | 2, 5 |
| N'Dama | NDAM | – | Dual purpose | Guinea (West Africa) | 1 |
| Norwegian Red | NRED | EU | Dairy | Norway | 1 |
| Piedmontese | PDMT | EU | Dual purpose | Italy | 1, 2 |
| Red Angus | RANG | BI/EU | Beef | Scotland | 1 |
| Red Poll | REDP | BI | Beef | England | 2 |
| Romagnola | ROMG | EU | Beef/draft | Italy | 1 |
| Scottish Highland | SCHL | BI | Beef | Scotland | 2 |
| Simmental | SIMM | EU | Dual purpose/draft | Switzerland | 2, 4 |
| South Devon | SDEV | BI | Beef | England | 2 |
| Sussex | SUSX | BI | Beef/draft | England | 2 |
| Welsh Black | WBLK | BI | Dual purpose | Wales | 2 |
| White Park | WHPK | BI | Dual purpose/draft | England | 2 |

¹Gibbs et al. (2009); ²Decker et al. (2009); ³Gautier et al. (2010); ⁴Matukumalli et al. (2009); ⁵Flori et al. (2009).

and genomic inbreeding, all SNPs genotyped (54,057) were filtered within breeds/populations as detailed above. However, an additional filtering procedure was used to remove SNPs deviating from Hardy-Weinberg equilibrium (HWE) with a *P*-value threshold of < 0.0001. Also, for the *N_e* analysis, a more stringent MAF threshold of 0.05 was used.

Generation of Identity-by-State (IBS) Matrix

Using the filtered genome-wide SNP data, PLINK v1.07 was also used to generate identity-by-state (IBS) values for all pairs of Kerry cattle (KY92 and KY12), including the nine blinded sample duplicates for sample verification and tracking purposes.

Construction of Phylogenetic Trees and Ancestry Graphs

Maximum likelihood (ML) phylogenetic trees with ancestry graphs were generated for the EU and BI data sets using the TreeMix (version 1.12) software package (Pickrell and Pritchard, 2012). The West African *B. taurus*

NDAM breed sample (*n* = 22) was used as an outgroup. TreeMix was run without using SNP blocks (as described in the TreeMix software documentation) and ML phylogenetic trees were generated with no migration edges (*m* = 0) up to ten migration edges (*m* = 10).

Population Differentiation and Genetic Structure

To visualise the main axes of genomic variation among cattle breeds and individual animals, multivariate principal component analysis (PCA) was performed for the composite EU and BI SNP data sets using SMARTPCA from the EIGENSOFT package (version 4.2) with default settings (Patterson et al., 2006).

To further investigate genetic structure and admixture history for Kerry cattle and other breeds the fastSTRUCTURE software package (Raj et al., 2014) was used to analyse the EU and BI data sets for a range of *K* possible ancestral populations (*K* = 2–15). For the present study, the simple prior approach described by Raj et al. (2014) was used, which is sufficient for

modelling population/breed divergence. To identify the “true” K -value for the number of ancestral populations, a series of fastSTRUCTURE runs with pre-defined K -values were examined using the *chooseK.py* script (Raj et al., 2014). Outputs from the fastSTRUCTURE analyses were visualised using the DISTRUCT software program (Rosenberg et al., 2002) using standard parameters.

Modelling Current and Historical Effective Population Size (N_e)

Current and historical N_e trends were modelled with genome-wide SNP linkage disequilibrium (LD) data for the KY92 and KY12 populations plus a selection of BI and EU breeds using the SNeP software tool as described by Barbato et al. (2015). This method facilitates estimation of historical N_e values from SNP linkage disequilibrium (LD) data using the following equation (Corbin et al., 2012):

$$N_{T(t)} = (4f(c_t))^{-1} \left(E[r_{adj}^2 | c_t]^{-1} - \alpha \right)$$

where N_T is the effective population size t generations ago calculated as $t = (2f(c_t))^{-1}$ (Hayes et al., 2003), c_t is the recombination rate defined for a specific physical distance between SNP markers, r_{adj}^2 is the LD value adjusted for sample size and the recommended default α value = 1 was used to correct for the occurrence of mutation (Barbato et al., 2015). In addition, the SNeP program option for unphased SNP data was used for the analyses described here.

Evaluation of Genomic Inbreeding and Runs of Homozygosity (ROH)

Individual animal genomic inbreeding was evaluated as genome-wide autozygosity estimated from SNP data using ROH and the F_{ROH} statistic introduced by McQuillan et al. (2008). The F_{ROH} statistic was calculated as the ratio of the total length of defined ROH (L_{ROH}) to the total length of the autosomal genome covered by SNPs:

$$F_{ROH} = \frac{\sum L_{ROH}}{L_{AUTO}}$$

PLINK v1.07 was used to define ROH using a sliding window approach and procedures modified from previous recommendations for Illumina® Bovine SNP50 BeadChip and similar SNP data sets (Purfield et al., 2012, 2017). The criteria for defining individual ROH were set such that the ROH was required to be at least 500 kb in length, with a minimum density of one SNP per 120 kb and that there was a gap of at least 1,000 kb between each ROH. A sliding window of 50 SNPs was incrementally advanced one SNP at a time along the genome; each discrete window could contain a maximum of one heterozygous SNP and no more than two SNPs with missing genotypes. Following Purfield et al. (2012) all filtered genomic SNPs (without a MAF threshold), including those located in centromeric regions, were used to estimate F_{ROH} values for individual animals.

Genome-Wide Detection of Signatures of Selection and Functional Enrichment Analysis

In the absence of hard selective sweeps, single selection tests using high-density SNP data do not perform well in detecting signatures of selection from individual livestock breeds (Kemper et al., 2014). Therefore, for the present study, genomic signatures of selection were identified using the composite selection signal (CSS) method introduced by Randhawa et al. (2014). The CSS method has been shown to be a robust and sensitive approach for detecting genomic signatures of selection underlying microevolution of complex traits in livestock (Randhawa et al., 2015). The CSS is a weighted index of signatures of selection from multiple estimates; it is a nonparametric procedure that uses fractional ranks of constituent tests and does not depend on assumptions about the distributions of individual test results.

As described in detail by Randhawa et al. (2014), the CSS method can be used to combine the fixation index (F_{ST}), the change in selected allele frequency (ΔSAF) and the cross-population extended haplotype homozygosity ($XP-EHH$) tests into one composite statistic for each SNP in a population genomics data set. For the present study, we used 36,621 genome-wide SNPs genotyped in 98 individual Kerry cattle samples (from both the KY92 and KY12 populations) and a sample of 102 randomly selected cattle (six random cattle from each breed of the EU data set). To mitigate against false positives, genomic selection signatures were only considered significant if at least one SNP from the set of the top 0.1% genome-wide CSS scores was flanked by at least five SNPs from the set of the top 1% CSS scores.

The Ensembl BioMart data mining resource (Smedley et al., 2015) was used to identify genes within ± 1.0 Mb of each selection peak (Ensembl release 90, August 2017). Following this, Ingenuity® Pathway Analysis (IPA®: Qiagen, Redwood City, CA, USA; release date June 2017) was used to perform an overrepresentation enrichment analysis (ORA) with this gene set to identify canonical pathways and functional processes of biological importance. The total gene content of Ensembl release 90 version of the UMD3.1 bovine genome assembly (Zimin et al., 2009) was used as the most appropriate reference gene set for these analyses (Timmons et al., 2015).

RESULTS AND DISCUSSION

Sample Removal and SNP Filtering and Quality Control

Genomic non-exclusion identified 20 parent-offspring pairs from the KY12 population and one sample from each pair was randomly removed (sample codes: KY12_01, KY12_05, KY12_13, KY12_14, KY12_17, KY12_18, KY12_19, KY12_46, KY12_55, KY12_67). Thereafter, general SNP quality control and filtering led to additional samples being excluded (KY12_26, KY12_28 and KY12_54), giving a total filtered KY12 population sample of 62 animals for downstream population genomics analyses.

After SNP quality control and filtering across the two composite data sets (EU and BI), there were 36,621 autosomal SNPs from 605 individual animals in the EU data set and there were 37,395 autosomal SNPs from 351 animals in the BI data set. When the West African NDAM breed sample ($n = 22$) was included for the ML phylogenetic tree and ancestry graph analyses, the number of SNPs used was 36,000 from 627 animals for the EU data set and 37,490 from 373 animals for the BI data set. The final numbers of SNPs used for individual breed/population analyses of N_e and genomic inbreeding after all quality control and filtering (including additional filtering for deviations from HWE) are shown in **Table 2**.

All data sets, including EU and BI composite data sets and individual breed/population data sets had total SNP call rates of $> 99\%$. The IBS values estimated for Kerry cattle (KY92 and KY12) from filtered genome-wide SNP data are reported in Supplementary Table 1 and described

further in section Genomic Relationship and Analysis of Inbreeding.

Observed Heterozygosity (H_o) Estimated from Genome-Wide SNP Data

Table 2 provides genome-wide H_o values for each of the breeds/populations used for the present study. The lowest genome-wide H_o value was observed for the West Africa NDAM *B. taurus* breed, which is likely a consequence of ascertainment bias introduced by a focus on polymorphic SNPs in European *B. taurus* during design of the Illumina® Bovine SNP50 BeadChip (Matukumalli et al., 2009).

Generally, as shown in **Table 2** for the EU and BI breeds and populations, local landrace or heritage breeds display lower H_o values compared to more widespread production breeds such as the Simmental (SIMM), Holstein (HOLS) or Charolais (CHAR) breeds. In addition, as might be expected, production breeds

TABLE 2 | Breed/population sample size, observed heterozygosity and SNP filtering information.

| Breed/population | Code | Data set | Sample size (n) post-filtering | Observed heterozygosity H_o | No. SNPs N_e modelling | No. SNPs genomic inbreeding |
|-----------------------|------|----------|---------------------------------------|----------------------------------|-----------------------------|--------------------------------|
| Angus | ANGU | BI/EU | 72 | 0.3048 | 31,413 | 39,576 |
| Belted Galloway | BGAL | BI | 4 | 0.2902 | 25,997 | 39,582 |
| British Shorthorn | BSHN | BI/EU | 10 | 0.2549 | 29,038 | 39,576 |
| Brown Swiss | BRSW | EU | 31 | 0.2894 | – | – |
| Charolais | CHAR | EU | 48 | 0.3209 | – | – |
| Devon | DEVN | BI | 4 | 0.2859 | – | – |
| Dexter | DXTR | BI | 4 | 0.2458 | 24,753 | 37,903 |
| English Longhorn | ELHN | BI | 3 | 0.2232 | – | – |
| Finnish Ayrshire | FAYR | BI/EU | 7 | 0.3064 | – | – |
| Galloway | GALL | BI | 4 | 0.2942 | – | – |
| Gelbvieh | GELB | EU | 8 | 0.3125 | – | – |
| Guernsey | GNSY | BI/EU | 19 | 0.2764 | – | 50,323 |
| Hereford | HRFD | BI/EU | 35 | 0.2964 | – | 39,535 |
| Holstein | HOLS | EU | 70 | 0.3192 | 36,152 | 43,135 |
| Jersey | JRSY | BI/EU | 44 | 0.2718 | 31,358 | 43,136 |
| Kerry sampled 1991/92 | KY92 | BI/EU | 36 | 0.2965 | 37,556 | 51,731 |
| Kerry sampled 2011/12 | KY12 | BI/EU | 62 | 0.3042 | 36,428 | 50,756 |
| Limousin | LIMS | EU | 45 | 0.3122 | – | – |
| Lincoln Red | LNCR | BI | 7 | 0.2789 | 26,350 | 34,173 |
| Montbeliarde | MONT | EU | 31 | 0.3019 | – | – |
| N'Dama | NDAM | – | 22 | 0.2158 | – | – |
| Norwegian Red | NRED | EU | 20 | 0.3190 | – | – |
| Piedmontese | PDMT | EU | 23 | 0.3240 | – | – |
| Red Angus | RANG | BI/EU | 14 | 0.3092 | – | – |
| Red Poll | REDP | BI | 5 | 0.2905 | – | – |
| Romagnola | ROMG | EU | 21 | 0.2943 | – | – |
| Scottish Highland | SCHL | BI | 8 | 0.2823 | – | – |
| Simmental | SIMM | EU | 9 | 0.3136 | – | – |
| South Devon | SDEV | BI | 3 | 0.3070 | – | – |
| Sussex | SUSX | BI | 4 | 0.2792 | – | – |
| Welsh Black | WBLK | BI | 2 | 0.3203 | – | – |
| White Park | WHPK | BI | 4 | 0.2270 | – | – |

originally derived from minor island populations (Jersey [JRSY] and Guernsey [GNSY]) also exhibit relatively low H_o values. In the context of genetic conservation it is therefore encouraging that the KY92 and KY12 population samples display intermediate H_o values that are at the upper end of the range observed for the heritage breeds.

Maximum Likelihood Phylogenetic Ancestry Graphs Using Genome-Wide SNP Data

To examine microevolutionary patterns of genetic differentiation and gene flow among cattle breeds and populations, ML phylogenetic ancestry graphs were generated using TreeMix. For the EU data set, the ML tree topology was consistent for all values of m , with the exception of $m = 2$ migration edges, where the Hereford breed (HRFD) was observed to group

with the HOLS breed. The ML tree generated with $m = 5$ is shown in **Figure 2**, which highlights the genetic similarity of the Northern European breeds (British, Irish and Scandinavian). As expected the two Kerry population samples (KY92 and KY12) are genetically very similar and emerge on the same branch as the HRFD breed. It is also noteworthy that there is a high-weight migration edge between the British Shorthorn breed (BSHN) and the root of the two Kerry population samples, supporting the hypothesis of historical gene flow from the British Shorthorn breed into the ancestral population of modern Kerry cattle (Curran, 1990).

For the ML trees generated using the BI data set, breed/population differentiation was less apparent, possibly due to similar biogeographical origins for these breeds and/or smaller sample sizes for some of the populations sampled. **Figure 3** shows the ML tree generated with $m = 5$ for the BI data set. For $m = 5$, all migration edges stem from the BSHN/Lincoln Red (LNCR)




FIGURE 2 | Maximum likelihood (ML) phylogenetic tree network graph with five migration edges ($m = 5$) generated for genome-wide SNP data (36,000 autosomal SNPs) from European cattle breeds (EU data set). The West African taurine N'Dama breed sampled in Guinea is included as a population outgroup. Coloured lines and arrows show migration edges that model gene flow between lineages with different migration weights represented by the colour gradient.




FIGURE 3 | Maximum likelihood (ML) phylogenetic tree network graph with five migration edges ($m = 5$) generated for genome-wide SNP data (37,490 autosomal SNPs) from cattle breeds of British and Irish origin (BI data set). The West African taurine N'Dama breed sampled in Guinea is included as a population outgroup. Coloured lines and arrows show migration edges that model gene flow between lineages with different migration weights represented by the colour gradient.

branch, including a medium-weight migration edge connecting to the Kerry cattle branch. These results support the hypothesis that there was significant gene flow during the eighteenth and nineteenth centuries from British Shorthorn cattle into the ancestral populations for a range of modern British and Irish cattle breeds (Grobet et al., 1998; Feliu et al., 2011, 2015).

Multivariate Principal Component Analysis of Genome-Wide SNP Data


To investigate inter- and intra-population genomic diversity and genetic relationship among individual animals from multiple cattle breeds and populations, PCA was performed using genome-wide SNP data. Principal component plots of the first (PC1) and second (PC2) principal components are shown in **Figures 4, 5** for the EU and BI data sets, respectively.

In **Figure 4**, for the EU data set, PC1 and PC2 account for 18.2% and 16.8% of the total variation for PC1–10, respectively. The PC1 plot axis differentiates the British Angus (ANGU), Red Angus (RANG) and BSHN and Irish KY92 and KY12 populations from the rest of the European breeds, including the British HRFD and GNSY and JRSY Channel Islands breeds. In addition, the ANGU and RANG and the Kerry (KY92 and KY12) emerge at the opposite extremes of the PC2 plot axis. In **Figure 5**, for the BI data set, PC1 and PC2 account for 23.7% and 22.7% of the total variation for PC1–10, respectively. The PC1 plot axis recapitulates PC2 in **Figure 4** and differentiates the Kerry (KY92 and KY12) from the ANGU and RANG breeds with the other British breeds emerging between these two extremes. These results highlight the genetic distinctiveness of the Kerry cattle

breed in comparison to a wide range of British production and heritage landrace cattle breeds and support their status as an important cattle genetic resource that should be prioritised for conservation.

The PC2 plot axis in **Figure 5** differentiates the HRFD breed from the other British and Irish breeds and reveals substantial genetic diversity among individual HRFD animals. However, in this context, it is important to note that the pattern of genetic diversity revealed here for the HRFD population sample may be due to ascertainment bias as a consequence of the strategy used to design the Illumina[®] Bovine SNP50 BeadChip. In this regard, many of the SNPs that constitute this first-generation SNP array were identified from heterozygous positions in the inbred Hereford female (L1 Dominette 01449) bovine genome assembly or through comparisons of random shotgun reads from six diverse cattle breeds that were aligned directly to the same Hereford genome assembly (Matukumalli et al., 2009). This approach to SNP array design will inevitably lead to elevated intrabreed genomic variation using the Illumina[®] Bovine SNP50 BeadChip with Hereford cattle (Meuwissen, 2009) and accounts for the dispersed pattern of individual HRFD samples in **Figure 5**.

Examination of **Figures 4, 5** indicates that two of the KY12 animals sampled may exhibit a genetic signature of ancestral crossbreeding with another cattle population, which, anecdotally, is likely to have been due to crossbreeding with Angus cattle. Therefore, another PCA plot was generated (Supplementary Figure 1) that shows PC1 and PC2 for individual animals from the KY92, KY12, ANGU and RANG population samples. The two






FIGURE 5 | Principal component analysis plot constructed for PC1 and PC2 from genome-wide SNP data (37,395 autosomal SNPs) for the BI data set of 351 individual animals. The smaller histogram plot shows the relative variance contributions for the first 10 PCs and PC1 and PC2 account for 23.7% and 22.7% of the total variation for PC1–10, respectively.

animals exhibiting a genetic signature of possible crossbreeding (KY12_06 and KY12_58) are indicated on Supplementary Figure 1. Notwithstanding the KY12_06 and KY12_58 data points, the genetic similarity among all Kerry cattle sampled is evident by comparison of the tight KY92 and KY12 sample cluster to the dispersion of the ANGU and RANG samples on the PCA plot in Supplementary Figure 1.

The values for the variation accounted for by PC3, PC4 and PC5 in **Figure 4** (EU data set) are relatively high (15.7, 13.8, and 10.4%, respectively). For **Figure 5** (BI data set), the variation accounted for by PC3 is also relatively high (19.1%). Therefore, we generated additional PCA plots of PC1 for each of the two data sets vs. these additional principal components (see Supplementary Figures 2–5).

Analysis of Genetic Structure Using Genome-Wide SNP Data

The results of the fastSTRUCTURE analyses using the EU and BI data sets are shown in **Figures 6, 7**, respectively. For both analyses, the Kerry cattle (KY92 and KY12) cluster as a single group at $K = 2$ and are differentiated from all other European or British and Irish cattle breeds. The other breed group that is clearly differentiated at $K = 2$ in **Figure 7** is the cluster composed of the ANGU and RANG breeds. These results mirror the pattern shown for PC1 in **Figure 5**, and again emphasise the genetic distinctiveness of Kerry cattle compared to other European production and landrace heritage breeds. Using the *chooseK.py* script the “true” number of clusters corresponding to

the likely number of ancestral populations was estimated to be between 12 and 14 for the EU data set and either 7 or 8 for the BI data set.

For both data sets, animals from the KY12 population sample appear to be more genetically homogenous compared to the KY92 population sample. This observation may be a consequence of increasing use, since the early 1990s, of small numbers of artificial insemination (AI) Kerry sires. It is also noteworthy that the two individual animals detected with a substantial signature of putative historical crossbreeding (KY12_06 and KY12_58) show marked patterns of population admixture in the fastSTRUCTURE results, which are indicated by red arrows in **Figures 6, 7**.

Modelling Historical Effective Population Size (N_e) Using Genome-Wide SNP Data

The results from modelling historical N_e in a selection of production and heritage cattle breeds and populations (KY92, KY12, DXTR, BSHN, BGAL, LNCR, ANGU, JRSY, and HOLS) are provided in Supplementary Table 2 and visualised in **Figure 8**. The “demographic fingerprints” (Barbato et al., 2015) of the two Kerry populations shown in **Figure 8** and tabulated in Supplementary Table 2 are more similar to those of the production breeds with large census populations (BSHN, ANGU, JRSY, HOLS) than the other heritage breeds with relatively small census population sizes (DXTR, BGAL, LNCR). The KY92, KY12, BSHN, ANGU, JRSY, and HOLS populations show a declining trend from historical N_e peaks between 1,500 and 2,000




FIGURE 6 | Hierarchical clustering of individual animals using genome-wide SNP data (36,621 autosomal SNPs) for the EU data set of 605 individual animals. Results are shown for modelled ancestral populations $K = 2-14$. The cluster numbers corresponding to the likely number of ancestral populations are highlighted with a light red overlay and the two outlier Kerry samples (KY12_06 and KY12_58) are indicated with red arrows.

more than 900 generations ago to N_e values estimated to be less than 200 within the last 20 generations. On the other hand, the DXTR, BSHN, BGAL and LNCR populations display a more severe decline from historical N_e peaks between 2,500 and 4,000 more than 900 generations ago to N_e values estimated to be less than 150 within the last 20 generations.

It is important to keep in mind these N_e trends may be partly a consequence of the relatively small sample sizes for the DXTR, BGAL, and LNCR breeds (see **Table 2**), coupled with different histories of migration, gene flow and, in particular, strong artificial selection in the production cattle populations. Notwithstanding these caveats, the most recent modelled N_e values for the KY92 and KY12 population samples are 89 and 88, respectively. These values are N_e estimates for 12 generations in the past and assuming a generation interval of between 4 and 6 years, which is based on a pedigree estimate from a similar heritage cattle population of 5.66 (Mészáros et al., 2015), this corresponds to between 48 and 72 years before 2012 (for the KY12 population). This is approximately the period between 1940 and 1965, which is during the time that the Kerry breed started to decline precipitously in census population size and also

N_e estimated from herd book data (O'hUigín and Cunningham, 1990; Food and Agriculture Organization, 2007).

From a conservation perspective, livestock populations generally exhibit N_e values relative to total census population sizes (N_c) that are substantially lower than seen in comparable wild mammal populations (Hall, 2016). Also, estimation of N_e using methods such as SNeP that leverage genome-wide SNP linkage disequilibrium (LD) data will tend to underestimate N_e because of physical linkage between many of the SNPs in the data set (Waples et al., 2016). Nevertheless, taking this into account, there is still cause for concern that the most recent N_e values modelled for the KY92 and KY12 population samples are below the critical N_e threshold of 100 recommended by Meuwissen (2009) for long-term viability of discrete livestock breeds and populations.

Genomic Relationship and Analysis of Inbreeding

Supplementary Table 1 shows a genomic relationship matrix in terms of genotype IBS for the genome-wide SNP data generated for individual animals in the KY92 and KY12 population





FIGURE 7 | Hierarchical clustering of individual animals using genome-wide SNP data (37,395 autosomal SNPs) for the BI data set of 351 individual animals. Results are shown for modelled ancestral populations $K = 2-9$. The cluster numbers corresponding to the likely number of ancestral populations are highlighted with a light red overlay and the two outlier Kerry samples (KY12_06 and KY12_58) are indicated with red arrows.

samples. Close genomic relationship between individual animals sampled from the same herd is evident in the SNP genotype IBS values between samples. In addition, the relatively low genomic relationship between the KY12_06 and KY12_58 outlier samples (Figures 4–7) and the rest of the Kerry cattle sampled is also evident in Supplementary Table 1. These data emphasise the value of intrapopulation genomic relationship values for identifying animals (in this case, KY12_06 and KY12_58) that should not be used in breeding programmes. They also highlight the potential of genome-wide SNP data for providing a systematic approach to prioritising males and females with minimum genomic relationship for breeding to minimize loss of genetic diversity and maintain or increase N_e (Gandini et al., 2004; Meuwissen, 2009; de Cara et al., 2011, 2013).

Genome-wide autozygosity estimated from SNP data using F_{ROH} and the F_{ROH} statistic are visualised in Figure 9 for individual animals from the KY92 and KY12 population samples and a range of European comparator breeds. Additional summary ROH data is provided in Supplementary Table 3 and also Supplementary Figure 6, which reveals marked

inter-population differences in ROH length and demonstrates that the SNP density of the Illumina® Bovine SNP50 BeadChip is too low to reliably capture ROH below 5 Mb in length, an observation previously reported by Purfield et al. (2012).

There is significant variation in F_{ROH} values among individual animals and between breeds and populations. The non-parametric Wilcoxon rank sum test was performed on F_{ROH} distributions for all pairwise population/breed comparisons with application of the Bonferroni correction P -value adjustment for multiple statistical tests (Supplementary Table 4). This analysis demonstrated that the KY12 population sample exhibited a significantly higher mean F_{ROH} value than the KY92 population sample (0.098 vs. 0.079; $P_{\text{adjust}} = 0.0081$). This is important from a conservation genetics perspective, indicating that genome-wide autozygosity, which is highly correlated with conventional pedigree-based estimates of inbreeding (F_{PED}) for cattle (Purfield et al., 2012; Ferencaković et al., 2013; Martikainen et al., 2017), has increased for the Kerry cattle population in the 20 years between sampling of the KY92 and KY12 populations.



The importance of understanding and quantifying genome-wide autozygosity for genetic conservation purposes has recently been highlighted through correlation of F_{ROH} with inbreeding depression for a range of production traits in domestic cattle (Bjelland et al., 2013; Pryce et al., 2014; Kim et al., 2015). Importantly, F_{ROH} has also been shown to correlate with inbreeding depression for bovine fertility traits in both males (Ferencaković et al., 2017) and females (Kim et al., 2015; Martikainen et al., 2017). Finally, according to basic population genetic principles, recent inbreeding captured by F_{ROH} will lead to recessive deleterious genomic variants emerging at a population level—a phenomenon that has been studied in both humans and cattle (Szpiech et al., 2013; Zhang et al., 2015).

Genome-Wide Signatures of Selection in the Kerry Cattle Breed

The results of the genome-wide scan for signatures of selection using the CSS method in the Kerry cattle breed are shown in **Figure 10**. Six distinct selection signatures were detected on BTA9, BTA12, BTA16, BTA17, BTA19, and BTA28. A total of 178 genes were located within the genomic ranges ± 1.0 Mb of selection peaks and 32 of these genes were located within the boundaries of a selection peak. Supplementary Table 5 provides detailed information for these 178 genes.

A single gene was located within the BTA9 selection peak—the phosphodiesterase 7B gene (*PDE7B*), which has been associated with neurobiological processes (de Gortari and Mengod, 2010) and has been previously linked to genetic changes associated with dog (*Canis lupus familiaris*) domestication and behaviour (Freedman et al., 2016). A single gene was also located

within the BTA16 selection peak—the dorsal inhibitory axon guidance protein gene (*DRAXIN*), which encodes a protein that regulates axon guidance, neural circuit formation and vertebrate brain development (Islam et al., 2009; Shinmyo et al., 2015). Twenty-four genes were located within the BTA17 selection peak, including *BICDL1*, *RAB35*, and *RNF10*, which have been associated with neurobiology and brain development (Hoshikawa et al., 2008; Schlager et al., 2010; Villarroel-Campos et al., 2016) and *SIRT4* and *COQ5* that function in cellular metabolism (Kawamukai, 2015; Elkhwanky and Hakkola, 2017). Six genes were located within the BTA28 selection peak, including, most notably, the Rho GTPase activating protein 22 gene (*ARHGAP22*), which has recently been associated with bovine fertility as an mRNA expression biomarker for oocyte competence in cumulus cells (Melo et al., 2017).

To obtain a broader perspective on natural and artificial selection acting at a population level on the Kerry cattle genome, a functional gene set enrichment approach (GSEA) was taken using IPA with the 178 genes located within ± 1.0 Mb of each selection peak (Supplementary Table 5). Of these 178 genes, 141 could be mapped to the IPA knowledgebase and the summary results for the IPA *Physiological System Development and Function* category are shown in Supplementary Table 6, revealing an enrichment of biological processes associated with nervous system development and behaviour. This functional enrichment coupled with the neurobiologically relevant single-gene selection peaks on BTA9 (*PDE7B*) and BTA16 (*DRAXIN*) suggests that natural and/or artificial selection related to brain development and behaviour has been important in the microevolution of the Kerry cattle breed. In this regard, it is therefore noteworthy




FIGURE 9 | Tukey box plots showing the distributions of F_{ROH} values estimated with genome-wide SNP data for the KY92 and KY12 populations and nine comparator heritage and production cattle breeds.

that Kerry cattle, including bulls, are recognised as being comparatively docile and easy to manage (Curran, 1990).

Genomics, Genetic Distinctiveness and Microevolution of Kerry Cattle: Implications for Breed Management and Genetic Conservation

The genome-wide phylogenetic and population genetic analyses detailed here demonstrate that Kerry cattle represent an important farm animal genetic resource, befitting the breed's status as a livestock population with a unique history of adaptation to the climate and physical geography of southwest Ireland at the edge of Western Europe. Notably, from a genetic conservation and breed management perspective, high-resolution comparative PCA (Figures 4, 5) and genetic clustering results (Figures 6, 7) demonstrate that Kerry cattle are markedly distinct from other British and European cattle populations. This observation may also be placed in the context of recent paleogenomic studies that have detected ancient gene flow from wild British aurochs (*B. primigenius*) into the ancestors of present-day Kerry cattle (Orlando, 2015; Park et al., 2015; Upadhyay et al., 2017).

The current genetic status of the Kerry cattle population is underlined by analyses of genetic effective population size (N_e) and inbreeding using genome-wide SNP data. As shown in Table 2, genome-wide observed heterozygosity (H_o) is relatively high in the KY92 and K12 population samples, particularly

for endangered heritage cattle breeds. However, it has been long recognised that monitoring N_e is a more important tool for rational breed management and long-term conservation of endangered livestock populations (Notter, 1999; Gandini et al., 2004; Biscarini et al., 2015). As shown in Figure 8 and Supplementary Table 2, the Kerry cattle population has a recent demographic trend of N_e decline, to the point where the most recent modelled N_e values are below the recommended threshold for sustainable breed management and conservation (Meuwissen, 2009). There is also cause for concern that genomic inbreeding estimated using genome-wide autozygosity (F_{ROH}) and visualised in Figure 9 has increased significantly in the 20-year period between the sampling of the KY92 and KY12 Kerry cattle populations.

In a more positive light, as shown in the present study, detection of discrete signatures of selection using the relatively low-density Illumina® Bovine SNP50 BeadChip is encouraging for wider studies of genome-wide microevolution in endangered heritage livestock populations such as Kerry cattle. Future surveys of heritage livestock populations that use higher-density SNP array platforms and ultimately whole-genome sequence data could provide exquisitely detailed information on the genomic regions and associated polygenic production, health, fertility and behavioural traits shaped, over many centuries, by the agro-ecology and pre-industrial farming systems of southwest Ireland.

In conclusion, the results presented here for the Kerry cattle population demonstrate that population genomics analyses of large SNP data sets can provide useful information concerning




FIGURE 10 | Manhattan plots of composite selection signal (CSS) results for Kerry cattle ($n = 98$) contrasted with EU cattle ($n = 102$). **(A)** Unsmoothed results. **(B)** Smoothed results obtained by averaging CSS of SNPs within each 1 Mb window. Red dotted line on each plot denotes the genome-wide 0.1% threshold for the empirical CSS scores. Red vertical arrows indicate selection peaks detected on BTA09, BTA12, BTA16, BTA17, BTA19, and BTA28.

the microevolution and recent genetic history of heritage livestock breeds. In particular, we would recommend that comparable surveys in other populations consider the use of genome-wide scans for signatures of selection, which can provide a functional genomics perspective on evolutionary adaptations to particular agricultural environments and production systems.

DATA ACCESSIBILITY

The Illumina® Bovine SNP50 BeadChip SNP genotype data for the Kerry cattle KY92 and KY12 population samples generated for this study are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.8fk81>.

ETHICS STATEMENT

With the exception of Kerry cattle sampled during 2011–12, all samples and data was obtained from previously published scientific studies. The re-use of these samples and data is consistent with the 3Rs principles on replacement, refinement and reduction of animals in research (www.nc3rs.org.uk/the-3rs). For the 2011–12 Kerry cattle, population owners' consent to sample DNA for research was obtained and individual owners conducted sampling of animals using non-invasive nasal swabs. In this regard, scientific animal protection in Ireland is subject

to European Union Directive 2010/63/EU, which states that the Directive does not apply to “practices not likely to cause pain, suffering, distress or lasting harm equivalent to, or higher than, that caused by the introduction of a needle in accordance with good veterinary practice.”

AUTHOR CONTRIBUTIONS

DEM, DAM, AF, and JK conceived and designed the project; DEM, IWR, DAM, AF, and JK organised sample collection and genotyping; SB, GM, IWR, DAM, SP, CC, IASR, and DEM performed the analyses; SB and DEM wrote the manuscript and all authors reviewed and approved the final manuscript.

FUNDING

This work was supported by Department of Agriculture, Food and the Marine (DAFM) funding under the Genetic Resources for Food and Agriculture scheme (grant no: 10/GR/06); an Investigator Programme Grant from Science Foundation Ireland (SFI/08/IN.1/B2038); a Research Stimulus Grant from DAFM (RSF 06 406); a European Union Framework 7 Project Grant (KBBE-211602-MACROSYS); the Brazilian Science Without Borders Programme (CAPES grant no. BEX-13070-13-4) and the UCD MSc Programme in Evolutionary Biology.

ACKNOWLEDGMENTS

The authors wish to express their gratitude to the Kerry Cattle Society (www.kerrycattle.ie) for facilitating and encouraging this project. In particular, we thank Rosemary and Jeremy Hill, Matthew English Hayden, Jeanne Hendrick and the Irish Cattle Breeding Federation (ICBF – www.icbf.com) for expert guidance and assistance with animal sourcing and DNA sampling. We are also grateful to Professor Dan Bradley (Smurfit Institute of Genetics, Trinity College Dublin) for access to DNA sample

archives. In addition, we thank all of the Kerry cattle owners who provided access to animals, samples and pedigree information. Finally, we thank Weatherbys Scientific for provision of SNP array genotyping services (www.weatherbysscscientific.com).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2018.00051/full#supplementary-material>

REFERENCES

- Barbato, M., Orozco-terWengel, P., Tapio, M., and Bruford, M. W. (2015). SNeP: a tool to estimate trends in recent effective population size trajectories using genome-wide SNP data. *Front. Genet.* 6:109. doi: 10.3389/fgene.2015.00109
- Ben Jemaa, S., Boussaha, M., Ben Mehdi, M., Lee, J. H., and Lee, S. H. (2015). Genome-wide insights into population structure and genetic history of Tunisian local cattle using the Illumina BovineSNP50 Beadchip. *BMC Genomics* 16:677. doi: 10.1186/s12864-015-1638-6
- Beynon, S. E., Slavov, G. T., Farré, M., Sunduimijid, B., Waddams, K., Davies, B., et al. (2015). Population structure and history of the Welsh sheep breeds determined by whole genome genotyping. *BMC Genet.* 16:65. doi: 10.1186/s12863-015-0216-x
- Biscarini, F., Nicolazzi, E. L., Stella, A., Boettcher, P. J., and Gandini, G. (2015). Challenges and opportunities in genetic improvement of local livestock breeds. *Front. Genet.* 6:33. doi: 10.3389/fgene.2015.00033
- Bjelland, D. W., Weigel, K. A., Vukasinovic, N., and Nkrumah, J. D. (2013). Evaluation of inbreeding depression in Holstein cattle using whole-genome SNP markers and alternative measures of genomic inbreeding. *J. Dairy Sci.* 96, 4697–4706. doi: 10.3168/jds.2012-6435
- Boettcher, P. J., Hoffmann, I., Baumung, R., Drucker, A. G., McManus, C., Berg, P., et al. (2015). Genetic resources and genomics for adaptation of livestock to climate change. *Front. Genet.* 5:461. doi: 10.3389/fgene.2014.00461
- Bray, T. C., Chikhi, L., Sheppy, A. J., and Bruford, M. W. (2009). The population genetic effects of ancestry and admixture in a subdivided cattle breed. *Anim. Genet.* 40, 393–400. doi: 10.1111/j.1365-2052.2009.01850.x
- Burren, A., Neuditschko, M., Signer-Hasler, H., Frischknecht, M., Reber, I., Menzi, F., et al. (2016). Genetic diversity analyses reveal first insights into breed-specific selection signatures within Swiss goat breeds. *Anim. Genet.* 47, 727–739. doi: 10.1111/age.12476
- Corbin, L. J., Liu, A. Y. H., Bishop, S. C., and Woolliams, J. A. (2012). Estimation of historical effective population size using linkage disequilibria with marker data. *J. Anim. Breed. Genet.* 129, 257–270. doi: 10.1111/j.1439-0388.2012.01003.x
- Curran, P. L. (1990). *Kerry and Dexter Cattle and other Ancient Irish Breeds, a History*. Dublin: Royal Dublin Society.
- de Cara, M. A., Fernández, J., Toro, M. A., and Villanueva, B. (2011). Using genome-wide information to minimize the loss of diversity in conservation programmes. *J. Anim. Breed. Genet.* 128, 456–464. doi: 10.1111/j.1439-0388.2011.00971.x
- de Cara, M. A., Villanueva, B., Toro, M. A., and Fernández, J. (2013). Using genomic tools to maintain diversity and fitness in conservation programmes. *Mol. Ecol.* 22, 6091–6099. doi: 10.1111/mec.12560
- Decker, J. E., Pires, J. C., Conant, G. C., McKay, S. D., Heaton, M. P., Chen, K., et al. (2009). Resolving the evolution of extant and extinct ruminants with high-throughput phylogenomics. *Proc. Natl. Acad. Sci. U.S.A.* 106, 18644–18649. doi: 10.1073/pnas.0904691106
- Decker, J. E., Taylor, J. F., Kantanen, J., Millbrooke, A., Schnabel, R. D., Alexander, L. J., et al. (2016). Origins of cattle on Chirikof Island, Alaska, elucidated from genome-wide SNP genotypes. *Heredity (Edinb)* 116, 502–505. doi: 10.1038/hdy.2016.7
- de Gortari, P., and Mengod, G. (2010). Dopamine D1, D2 and mu-opioid receptors are co-expressed with adenylyl cyclase 5 and phosphodiesterase 7B mRNAs in striatal rat cells. *Brain Res.* 1310, 37–45. doi: 10.1016/j.brainres.2009.11.009
- Department of Agriculture Food and the Marine (2014). *State of Biodiversity for Food and Agriculture in Ireland*. Dublin: Department of Agriculture, Food and the Marine.
- Department of Agriculture Food and the Marine (2017). *Kerry Cattle Premium Scheme: Scheme Objectives and Terms and Conditions*. Dublin: Department of Agriculture, Food and the Marine.
- Elkhwanky, M. S., and Hakkola, J. (2017). Extranuclear sirtuins and metabolic stress. *Antioxid. Redox Signal.* doi: 10.1089/ars.2017.7270. [Epub ahead of print].
- Felius, M., Koolmees, P. A., Theunissen, B., Consortium, E. C. G. D., and Lenstra, J. A. (2011). On the breeds of cattle—historic and current classifications. *Diversity (Basel)* 3, 660–692. doi: 10.3390/d3040660
- Felius, M., Theunissen, B., and Lenstra, J. A. (2015). Conservation of cattle genetic resources: the role of breeds. *J. Agric. Sci.* 153, 152–162. doi: 10.1017/S0021859614000124
- Ferenčaković, M., Hamzić, E., Gredler, B., Solberg, T. R., Klemetsdal, G., Curik, I., et al. (2013). Estimates of autozygosity derived from runs of homozygosity: empirical evidence from selected cattle populations. *J. Anim. Breed. Genet.* 130, 286–293. doi: 10.1111/jbg.12012
- Ferenčaković, M., Sölkner, J., Kapš, M., and Curik, I. (2017). Genome-wide mapping and estimation of inbreeding depression of semen quality traits in a cattle population. *J. Dairy Sci.* 100, 4721–4730. doi: 10.3168/jds.2016-12164
- Flori, L., Fritz, S., Jaffrézic, F., Boussaha, M., Gut, I., Heath, S., et al. (2009). The genome response to artificial selection: a case study in dairy cattle. *PLoS ONE* 4:e6595. doi: 10.1371/journal.pone.0006595
- Food and Agriculture Organization (2007). *The State of the World's Animal Genetic Resources for Food and Agriculture*. Rome: FAO Commission on Genetic Resources for Food and Agriculture Assessments.
- Food and Agriculture Organization (2015). *The Second Report on the State of the World's Animal Genetic Resources for Food and Agriculture*. Rome: FAO Commission on Genetic Resources for Food and Agriculture Assessments.
- Food and Agriculture Organization (2017). *Domestic Animal Diversity Information System (DAD-IS)*. Available online at: www.fao.org/dad-is/en (Accessed 01/05/2017).
- François, L., Wijnrocx, K., Colinet, F. G., Gengler, N., Hulsege, B., Windig, J. J., et al. (2017). Genomics of a revived breed: case study of the Belgian Campine cattle. *PLoS ONE* 12:e0175916. doi: 10.1371/journal.pone.0175916
- Freedman, A. H., Schweizer, R. M., Ortega-Del Vecchyo, D., Han, E., Davis, B. W., Gronau, I., et al. (2016). Demographically-based evaluation of genomic regions under selection in domestic dogs. *PLoS Genet.* 12:e1005851. doi: 10.1371/journal.pgen.1005851
- Gandini, G. C., Ollivier, L., Danell, B., Distl, O., Georgoudis, A., Groeneveld, E., et al. (2004). Criteria to assess the degree of endangerment of livestock breeds in Europe. *Livestock Prod. Sci.* 91, 173–182. doi: 10.1016/j.livprodsci.2004.08.001
- Gao, Y., Sun, J., Li, F., He, S., Sandven, S., Yan, Q., et al. (2015). Arctic sea ice and Eurasian climate: a review. *Adv. Atmos. Sci.* 32, 92–114. doi: 10.1007/s00376-014-0009-6
- Gautier, M., Laloë, D., and Moazami-Goudarzi, K. (2010). Insights into the genetic history of French cattle from dense SNP data on 47 worldwide breeds. *PLoS ONE* 5:e13038. doi: 10.1371/journal.pone.0013038
- Gibbs, R. A., Taylor, J. F., Van Tassell, C. P., Barendse, W., Eversole, K. A., Gill, C. A., et al. (2009). Genome-wide survey of SNP variation uncovers the genetic structure of cattle breeds. *Science* 324, 528–532. doi: 10.1126/science.1167936

- Grobet, L., Poncelet, D., Royo, L. J., Brouwers, B., Pirottin, D., Michaux, C., et al. (1998). Molecular definition of an allelic series of mutations disrupting the myostatin function and causing double-muscling in cattle. *Mamm. Genome* 9, 210–213. doi: 10.1007/s003359900727
- Groeneveld, L. F., Gregusson, S., Guldbrandtsen, B., Hiemstra, S. J., Hveem, K., Kantanen, J., et al. (2016). Domesticated animal biobanking: land of opportunity. *PLoS Biol.* 14:e1002523. doi: 10.1371/journal.pbio.1002523
- Hall, S. J. (2016). Effective population sizes in cattle, sheep, horses, pigs and goats estimated from census and herdbook data. *Animal* 10, 1778–1785. doi: 10.1017/S1751731116000914
- Hayes, B. J., Visscher, P. M., McPartlan, H. C., and Goddard, M. E. (2003). Novel multilocus measure of linkage disequilibrium to estimate past effective population size. *Genome Res.* 13, 635–643. doi: 10.1101/gr.387103
- Hill, W. G. (2014). Applications of population genetics to animal breeding, from Wright, Fisher and Lush to genomic prediction. *Genetics* 196, 1–16. doi: 10.1534/genetics.112.147850
- Hoffmann, I. (2010). Climate change and the characterization, breeding and conservation of animal genetic resources. *Anim. Genet.* 41(Suppl. 1), 32–46. doi: 10.1111/j.1365-2052.2010.02043.x
- Hoshikawa, S., Ogata, T., Fujiwara, S., Nakamura, K., and Tanaka, S. (2008). A novel function of RING finger protein 10 in transcriptional regulation of the myelin-associated glycoprotein gene and myelin formation in Schwann cells. *PLoS ONE* 3:e3464. doi: 10.1371/journal.pone.0003464
- Islam, S. M., Shimmyo, Y., Okafuji, T., Su, Y., Naser, I. B., Ahmed, G., et al. (2009). Draxin, a repulsive guidance protein for spinal cord and forebrain commissures. *Science* 323, 388–393. doi: 10.1126/science.1165187
- Iso-Touru, T., Tapio, M., Vilkkilä, J., Kiseleva, T., Ammosov, I., Ivanova, Z., et al. (2016). Genetic diversity and genomic signatures of selection among cattle breeds from Siberia, eastern and northern Europe. *Anim. Genet.* 47, 647–657. doi: 10.1111/age.12473
- Kantanen, J., Løvendahl, P., Strandberg, E., Eythorsdóttir, E., Li, M. H., Kettunen-Präbel, A., et al. (2015). Utilization of farm animal genetic resources in a changing agro-ecological environment in the Nordic countries. *Front. Genet.* 6:52. doi: 10.3389/fgene.2015.00052
- Kawamukai, M. (2015). Biosynthesis of coenzyme Q in eukaryotes. *Biosci. Biotechnol. Biochem.* 80, 23–33. doi: 10.1080/09168451.2015.1065172
- Kemper, K. E., Saxton, S. J., Bolormaa, S., Hayes, B. J., and Goddard, M. E. (2014). Selection for complex traits leaves little or no classic signatures of selection. *BMC Genomics* 15:246. doi: 10.1186/1471-2164-15-246
- Kim, E. S., Sonstegard, T. S., Van Tassell, C. P., Wiggans, G., and Rothschild, M. F. (2015). The relationship between runs of homozygosity and inbreeding in Jersey cattle under selection. *PLoS ONE* 10:e0129967. doi: 10.1371/journal.pone.0129967
- Kristensen, T. N., Hoffmann, A. A., Pertoldi, C., and Stronen, A. V. (2015). What can livestock breeders learn from conservation genetics and vice versa? *Front. Genet.* 6:38. doi: 10.3389/fgene.2015.00038
- Larson, G., and Fuller, D. Q. (2014). The evolution of animal domestication. *Annu. Rev. Ecol. Syst.* 45, 115–136. doi: 10.1146/annurev-ecolsys-110512-135813
- Larson, G., Piperno, D. R., Allaby, R. G., Purugganan, M. D., Andersson, L., Arroyo-Kalin, M., et al. (2014). Current perspectives and the future of domestication studies. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6139–6146. doi: 10.1073/pnas.1323964111
- MacHugh, D. E., Larson, G., and Orlando, L. (2017). Taming the past: ancient DNA and the study of animal domestication. *Annu. Rev. Anim. Biosci.* 5, 329–351. doi: 10.1146/annurev-animal-022516-022747
- MacHugh, D. E., Loftus, R. T., Cunningham, P., and Bradley, D. G. (1998). Genetic structure of seven European cattle breeds assessed using 20 microsatellite markers. *Anim. Genet.* 29, 333–340. doi: 10.1046/j.1365-2052.1998.295330.x
- MacHugh, D. E., Shriver, M. D., Loftus, R. T., Cunningham, P., and Bradley, D. G. (1997). Microsatellite DNA variation and the evolution, domestication and phylogeography of taurine and zebu cattle (*Bos taurus* and *Bos indicus*). *Genetics* 146, 1071–1086.
- MacHugh, D. E., Troy, C. S., McCormick, F., Olsaker, I., Eythorsdóttir, E., and Bradley, D. G. (1999). Early medieval cattle remains from a Scandinavian settlement in Dublin: genetic analysis and comparison with extant breeds. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 354, 99–108. doi: 10.1098/rstb.1999.0363
- Manunza, A., Cardoso, T. F., Noce, A., Martínez, A., Pons, A., Bermejo, L. A., et al. (2016). Population structure of eleven Spanish ovine breeds and detection of selective sweeps with BayeScan and hapFLK. *Sci. Rep.* 6:27296. doi: 10.1038/srep27296
- Martikainen, K., Tyrisevä, A. M., Matilainen, K., Pösö, J., and Uimari, P. (2017). Estimation of inbreeding depression on female fertility in the Finnish Ayrshire population. *J. Anim. Breed. Genet.* 134, 383–392. doi: 10.1111/jbg.12285
- Mastrangelo, S., Tolone, M., Di Gerlando, R., Fontanesi, L., Sardina, M. T., and Portolano, B. (2016). Genomic inbreeding estimation in small populations: evaluation of runs of homozygosity in three local dairy cattle breeds. *Animal* 10, 746–754. doi: 10.1017/S1751731115002943
- Matukumalli, L. K., Lawley, C. T., Schnabel, R. D., Taylor, J. F., Allan, M. F., Heaton, M. P., et al. (2009). Development and characterization of a high density SNP genotyping assay for cattle. *PLoS ONE* 4:e5350. doi: 10.1371/journal.pone.0005350
- McParland, S. (2013). *National Genetic Conservation Strategy Document*. Available online at: <https://www.agriculture.gov.ie>
- McQuillan, R., Leutenegger, A. L., Abdel-Rahman, R., Franklin, C. S., Pericic, M., Barac-Lauc, L., et al. (2008). Runs of homozygosity in European populations. *Am. J. Hum. Genet.* 83, 359–372. doi: 10.1016/j.ajhg.2008.08.007
- Melo, E. O., Cordeiro, D. M., Pellegrino, R., Wei, Z., Daye, Z. J., Nishimura, R. C., et al. (2017). Identification of molecular markers for oocyte competence in bovine cumulus cells. *Anim. Genet.* 48, 19–29. doi: 10.1111/age.12496
- Mészáros, G., Boison, S. A., Pérez O'Brien, A. M., Ferencaković, M., Curik, I., Da Silva, M. V., et al. (2015). Genomic analysis for managing small and endangered populations: a case study in Tyrol Grey cattle. *Front. Genet.* 6:173. doi: 10.3389/fgene.2015.00173
- Meuwissen, T. (2009). Genetic management of small populations: a review. *Acta Agric. Scandinavica Section A Anim. Sci.* 59, 71–79. doi: 10.1080/09064700903118148
- Notter, D. R. (1999). The importance of genetic diversity in livestock populations of the future. *J. Anim. Sci.* 77, 61–69. doi: 10.2527/1999.77161x
- O'hUigín, C., and Cunningham, E. P. (1990). Analysis of breeding structure of the Kerry breed. *J. Anim. Breed. Genet.* 107, 452–457. doi: 10.1111/j.1439-0388.1990.tb00057.x
- Orlando, L. (2015). The first aurochs genome reveals the breeding history of British and European cattle. *Genome Biol.* 16:225. doi: 10.1186/s13059-015-0793-z
- Park, S. D., Magee, D. A., McGettigan, P. A., Teasdale, M. D., Edwards, C. J., Lohan, A. J., et al. (2015). Genome sequencing of the extinct Eurasian wild aurochs, *Bos primigenius*, illuminates the phylogeography and evolution of cattle. *Genome Biol.* 16:234. doi: 10.1186/s13059-015-0790-2
- Patterson, N., Price, A. L., and Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genet.* 2:e190. doi: 10.1371/journal.pgen.0020190
- Pertoldi, C., Purfield, D. C., Berg, P., Jensen, T. H., Bach, O. S., Vingborg, R., et al. (2014). Genetic characterization of a herd of the endangered Danish Jutland cattle. *J. Anim. Sci.* 92, 2372–2376. doi: 10.2527/jas.2013-7206
- Phocas, F., Belloc, C., Bidanel, J., Delaby, L., Dourmad, J. Y., Dumont, B., et al. (2016a). Review: towards the agroecological management of ruminants, pigs and poultry through the development of sustainable breeding programmes. II. Breeding strategies. *Animal* 10, 1760–1769. doi: 10.1017/S1751731116001051
- Phocas, F., Belloc, C., Bidanel, J., Delaby, L., Dourmad, J. Y., Dumont, B., et al. (2016b). Review: towards the agroecological management of ruminants, pigs and poultry through the development of sustainable breeding programmes: I-selection goals and criteria. *Animal* 10, 1749–1759. doi: 10.1017/S1751731116000926
- Pickrell, J. K., and Pritchard, J. K. (2012). Inference of population splits and mixtures from genome-wide allele frequency data. *PLoS Genet.* 8:e1002967. doi: 10.1371/journal.pgen.1002967
- Pryce, J. E., Haile-Mariam, M., Goddard, M. E., and Hayes, B. J. (2014). Identification of genomic regions associated with inbreeding depression in Holstein and Jersey dairy cattle. *Genet. Sel. Evol.* 46:71. doi: 10.1186/s12711-014-0071-7
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., et al. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81, 559–575. doi: 10.1086/519795
- Purfield, D. C., Berry, D. P., McParland, S., and Bradley, D. G. (2012). Runs of homozygosity and population history in cattle. *BMC Genet.* 13:70. doi: 10.1186/1471-2156-13-70

- Purfield, D. C., McParland, S., Wall, E., and Berry, D. P. (2017). The distribution of runs of homozygosity and selection signatures in six commercial meat sheep breeds. *PLoS ONE* 12:e0176780. doi: 10.1371/journal.pone.0176780
- Raj, A., Stephens, M., and Pritchard, J. K. (2014). fastSTRUCTURE: variational inference of population structure in large SNP data sets. *Genetics* 197, 573–589. doi: 10.1534/genetics.114.164350
- Randhawa, I. A., Khatkar, M. S., Thomson, P. C., and Raadsma, H. W. (2014). Composite selection signals can localize the trait specific genomic regions in multi-breed populations of cattle and sheep. *BMC Genet.* 15:34. doi: 10.1186/1471-2156-15-34
- Randhawa, I. A., Khatkar, M. S., Thomson, P. C., and Raadsma, H. W. (2015). Composite selection signals for complex traits exemplified through bovine stature using multibreed cohorts of European and African *Bos taurus*. *G3 (Bethesda)* 5, 1391–1401. doi: 10.1534/g3.115.017772
- Rosenberg, N. A., Pritchard, J. K., Weber, J. L., Cann, H. M., Kidd, K. K., Zhivotovsky, L. A., et al. (2002). Genetic structure of human populations. *Science* 298, 2381–2385. doi: 10.1126/science.1078311
- Schlager, M. A., Kapitein, L. C., Grigoriev, I., Burzynski, G. M., Wulf, P. S., Keijzer, N., et al. (2010). Pericentrosomal targeting of Rab6 secretory vesicles by Bicaudal-D-related protein 1 (BICDR-1) regulates neuritogenesis. *EMBO J.* 29, 1637–1651. doi: 10.1038/emboj.2010.51
- Shinmyo, Y., Asrafuzzaman Riyadh, M., Ahmed, G., Bin Naser, I., Hossain, M., Takebayashi, H., et al. (2015). Draxin from neocortical neurons controls the guidance of thalamocortical projections into the neocortex. *Nat. Commun.* 6:10232. doi: 10.1038/ncomms10232
- Smedley, D., Haider, S., Durinck, S., Pandini, L., Provero, P., Allen, J., et al. (2015). The BioMart community portal: an innovative alternative to large, centralized data repositories. *Nucleic Acids Res.* 43, W589–W598. doi: 10.1093/nar/gkv350
- Smith, P., and Gregory, P. J. (2013). Climate change and sustainable food production. *Proc. Nutr. Soc.* 72, 21–28. doi: 10.1017/S0029665112002832
- Szpiech, Z. A., Xu, J., Pemberton, T. J., Peng, W., Zöllner, S., Rosenberg, N. A., et al. (2013). Long runs of homozygosity are enriched for deleterious variation. *Am. J. Hum. Genet.* 93, 90–102. doi: 10.1016/j.ajhg.2013.05.003
- Timmons, J. A., Szkop, K. J., and Gallagher, I. J. (2015). Multiple sources of bias confound functional enrichment analysis of global -omics data. *Genome Biol.* 16:186. doi: 10.1186/s13059-015-0761-7
- Troy, C. S., MacHugh, D. E., Bailey, J. F., Magee, D. A., Loftus, R. T., Cunningham, P., et al. (2001). Genetic evidence for Near-Eastern origins of European cattle. *Nature* 410, 1088–1091. doi: 10.1038/35074088
- Upadhyay, M. R., Chen, W., Lenstra, J. A., Goderie, C. R., MacHugh, D. E., Park, S. D., et al. (2017). Genetic origin, admixture and population history of aurochs (*Bos primigenius*) and primitive European cattle. *Heredity (Edinb)* 118, 169–176. doi: 10.1038/hdy.2016.79
- Vihma, T. (2014). Effects of Arctic sea ice decline on weather and climate: a review. *Surveys Geophys.* 35, 1175–1214. doi: 10.1007/s10712-014-9284-0
- Villaruel-Campos, D., Henríquez, D. R., Bodaleo, F. J., Oguchi, M. E., Bronfman, F. C., Fukuda, M., et al. (2016). Rab35 functions in axon elongation are regulated by P53-related protein kinase in a mechanism that involves Rab35 protein degradation and the microtubule-associated protein 1B. *J. Neurosci.* 36, 7298–7313. doi: 10.1523/JNEUROSCI.4064-15.2016
- Visser, C., Lashmar, S. F., Van Marle-Köster, E., Poli, M. A., and Allain, D. (2016). Genetic diversity and population structure in South African, French and Argentinian Angora goats from genome-wide SNP data. *PLoS ONE* 11:e0154353. doi: 10.1371/journal.pone.0154353
- Waples, R. K., Larson, W. A., and Waples, R. S. (2016). Estimating contemporary effective population size in non-model species using linkage disequilibrium across thousands of loci. *Heredity (Edinb)* 117, 233–240. doi: 10.1038/hdy.2016.60
- Wheeler, T., and von Braun, J. (2013). Climate change impacts on global food security. *Science* 341, 508–513. doi: 10.1126/science.1239402
- Williams, J. L., Hall, S. J., Del Corvo, M., Ballingall, K. T., Colli, L., Ajmone Marsan, P., et al. (2016). Inbreeding and purging at the genomic level: the Chillingham cattle reveal extensive, non-random SNP heterozygosity. *Anim. Genet.* 47, 19–27. doi: 10.1111/age.12376
- Zhang, Q., Guldbrandsen, B., Bosse, M., Lund, M. S., and Sahana, G. (2015). Runs of homozygosity and distribution of functional variants in the cattle genome. *BMC Genomics* 16:542. doi: 10.1186/s12864-015-1715-x
- Zimin, A. V., Delcher, A. L., Florea, L., Kelley, D. R., Schatz, M. C., Puiu, D., et al. (2009). A whole-genome assembly of the domestic cow, *Bos taurus*. *Genome Biol.* 10:R42. doi: 10.1186/gb-2009-10-4-r42

Conflict of Interest Statement: The authors IWR and SP are employed by IdentiGEN, Ltd.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Browett, McHugo, Richardson, Magee, Park, Fahey, Kearney, Correia, Randhawa and MacHugh. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.